



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 9/24	A1	(11) International Publication Number: WO 99/47128 (43) International Publication Date: 23 September 1999 (23.09.99)
(21) International Application Number: PCT/US99/05233 (22) International Filing Date: 10 March 1999 (10.03.99) (30) Priority Data: 09/044,446 19 March 1998 (19.03.98) US (71) Applicant: BRISTOL-MYERS SQUIBB COMPANY [US/US]; P.O. Box 4000, Princeton, NJ 08543-4000 (US). (72) Inventors: TIMMINS, Peter; 5 Heathbank Avenue, Irby, Merseyside L61 4XD (GB). DENNIS, Andrew, B.; 7 Pear Tree Close, Barnston, Merseyside L60 1YD (GB). VYAS, Kiren, A.; 34 Adisham Green, Sittingbourne, Kent ME10 2SR (GB). (74) Agents: RODNEY, Burton et al.; Bristol-Myers Squibb Com- pany, P.O. Box 4000, Princeton, NJ 08543-4000 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW. ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: BIPHASIC CONTROLLED RELEASE DELIVERY SYSTEM FOR HIGH SOLUBILITY PHARMACEUTICALS AND METHOD (57) Abstract <p>A biphasic controlled release delivery system for pharmaceuticals which have high water solubility, such as the antidiabetic metformin HCl salt, is provided which provides a dosage form that has prolonged gastric residence and includes (1) an inner solid particulate phase formed of substantially uniform granules containing a pharmaceutical having a high water solubility, and one or more hydrophilic polymers, one or more hydrophobic polymers and/or one or more hydrophobic materials such as one or more waxes, fatty alcohols and/or fatty acid esters, and (2) an outer solid continuous phase in which the above granules of inner solid particulate phase are embedded and dispersed throughout, the outer solid continuous phase including one or more hydrophilic polymers, one or more hydrophobic polymers and/or one or more hydrophobic materials such as one or more waxes, fatty alcohols and/or fatty acid esters, which may be compressed into tablets or filled into capsules. Methods for forming the so-described biphasic controlled release delivery system and using such biphasic controlled release delivery system for treating diabetes are also provided.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

BIPHASIC CONTROLLED RELEASE DELIVERY SYSTEM
FOR HIGH SOLUBILITY PHARMACEUTICALS AND METHOD

Field of the Invention

5 The present invention relates to a new dosage form
for highly water soluble medicaments, such as the
antidiabetic metformin, which provides for extended release
of the drug and also for prolonged gastric residence which
enables efficient delivery of drugs normally absorbed in
10 the upper gastrointestinal tract, and to a method for
preparing such dosage form.

Background of the Invention

15 Metformin is an antihyperglycemic agent of the
biguanide class used in the treatment of non-insulin
dependent diabetes mellitus (NIDDM). It is usually
marketed in the form of its hydrochloride salt as
Glucophage® (TM-BMS).

20 Metformin hydrochloride has intrinsically poor
permeability in the lower portion of the gastrointestinal
tract leading to absorption almost exclusively in the upper
part of the gastrointestinal tract. Its oral
bioavailability is in the range of 40 to 60% decreasing
with increasing dosage which suggests some kind of
25 saturable absorption process, or permeability/transit time
limited absorption. It also has a very high water
solubility (>300 mg/ml at 25°C). This can lead to
difficulty in providing a slow release rate from a
formulation and problems in controlling the initial burst
30 of drug from such a formulation. These two difficulties
are further compounded by the high unit dose, 500 mg per
tablet, usually required for metformin hydrochloride (1997-
PDR).

35 Drugs that have absorption limited to the upper
gastrointestinal tract coupled with poor absorption in the
distal small intestine, large intestine and colon are
usually regarded as inappropriate candidates for

formulation into oral controlled delivery systems. This limitation on absorption (for example, in the upper gastrointestinal tract) is referred to as the "absorption window".

5 The gastrointestinal tract functions to propel ingested material from the stomach (where digestion takes place) into the small intestine (where absorption principally occurs) and on to the large intestine (where water is absorbed/secreted as part of body fluid regulation
10 processes). Residence time for non-digestible materials in the stomach depends on whether one is dealing with a fed or a fasted subject. Typical gastric emptying times for particulate material (greater than a few millimeters in diameter) varies from a few tens of minutes in the fasted
15 state to a few hours in the fed state. Transit times through the small intestine are consistently of the order of 3 to 4 hours.

 Oral controlled release delivery systems function by releasing their payload of drug over an extended period of
20 time following administration. Thus, controlled release dosage forms may only spend a relatively short period in the regions of the gastrointestinal tract where good absorption of certain drugs can occur. The dosage form will pass on to regions of the intestine where absorption
25 of certain drugs is poor or non-existent, still releasing its contained drug albeit with a significant percentage of its payload still to be delivered. Drug when released from the dosage form in the circumstances described will not be absorbed. Thus, administration of a drug subject to a
30 window of absorption in a conventional controlled release delivery system can lead to subtherapeutic blood levels and ineffective treatment of the disease state for which the drug was intended.

 Drugs with very high solubility in water (for
35 example, greater than 100 mg/ml) can be difficult to formulate into a controlled release oral dosage form. Solubility is a driving force for a drug substance to

dissolve in water; the greater the solubility the greater the rate of dissolution when all other factors are maintained constant.

In a controlled release dosage form, the formulator
5 tries to reduce the rate of dissolution by, for example, embedding the drug in a polymeric matrix or surrounding it with a polymeric barrier membrane through which drug must diffuse to be released for absorption. To reduce the rate of release of drug from the dosage form to an appropriate
10 level consistent with the blood level profile desired for a drug possessing very high water solubility, very large amounts of polymer would be required for the matrix or barrier membrane. If the total daily dose of drug to be delivered is of the order of only a few milligrams this may
15 be feasible, but many drugs having the solubility properties described require total daily doses of the order of many hundreds of milligrams. Whilst it is possible to create oral controlled release dosage forms for such products by use of large amounts of polymer, an
20 unacceptably large dosage form may result.

A further problem with highly water soluble drugs formulated into a controlled release dosage form is that a significant and variable "burst" of drug can occur from these systems. The burst of highly water soluble drug is
25 the initial rapid release of drug that occurs from oral controlled release dosage forms when first contacting fluid, such as gastric fluids, prior to release controlling mechanisms of the dosage form establishing themselves and a stable release rate being provided. Hydration of any
30 polymer matrix used to formulate the dosage form is a pre-requirement of establishing a stable release rate. Thus, a readily hydrating polymer is required to establish the desired stable release rate. However, if the polymer used is slow to hydrate, then an undesirable variable burst can
35 occur.

Studies by Vidon et al (1) strongly suggest that there is permeability limited absorption of metformin.

Perfusing drug into the jejunum via an intubation technique showed a 2.5 fold greater area under the plasma concentration-time profile (a measure of the quantity of drug absorbed) compared with similar introduction of drug into the ileum. Drug was not detectable in plasma when drug was perfused into the colon. Drug will transit down the small intestine following dissolution from an ingested dosage form and, if absorption rate is slow, it is possible that drug can reach regions of poor permeability before absorption of a given dose is complete. In such a case, increasing the given dose may be predicted to result in a reduction in the percentage of administered dose absorbed.

Improvements in the therapeutic regimes employing metformin might be achieved by a dosage form that allows a reduction in dosing frequency, providing patient convenience that would probably improve compliance. Conventional extended release formulations have been demonstrated to invariably compromise the availability of metformin (2). This is probably because the dosage form carries a significant proportion of the drug content remaining to be released, as the dosage form is carried to regions of the gastrointestinal tract with very poor permeability to the drug. To reduce dosing frequency, the rate of release from the dosage form must be such as to extend effective plasma levels, but the potential for effective delivery at this rate is compromised by the combined influences of the significant reduction in permeability to the drug in passing from the proximal small intestine down to the colon and the limited residence time in the regions of the gastrointestinal tract where the drug is well absorbed. That transit time down the "useful" region of the gastrointestinal tract is only likely to be of the order of a few hours.

Maintained or even improved bioavailability from an extended release dosage form that releases metformin at a rate likely to provide the desired plasma levels of drug for an extended time period might, however, be possible

from a dosage form that has extended residence time in the upper gastrointestinal tract, resisting mechanisms that promote normal transit time for solid materials. That this principle might work in practice was demonstrated in an in-house study where metformin was co-administered with propantheline, an agent that reduces gastrointestinal motility. Compared with giving metformin alone, the combination provided an increased AUC, a delayed tmax and an extended time period over which therapeutically beneficial plasma levels of drug were maintained.

Giving a drug such as metformin for the treatment of diabetes with a further drug, such as propantheline, not used for the treatment of diabetes and where the sole intent of using the second agent is to achieve extended residence time in the upper GI tract, has many disadvantages although it is likely to allow effective extended delivery of metformin to an optimal absorption site. The co-administered drug may have other undesirable pharmacological effects or side effects deleterious to the patients well being and detract from the improved quality of life offered by the treatment for their diabetes. Furthermore, it may be difficult or impossible to appropriately co-formulate the two agents due to chemical compatibility issues or solubility differences, the latter preventing the required release rate of agent influencing residence time in the upper GI tract. Thus, the patient could be required to take separate, multiple medications to achieve the desired effect. The timing of taking the two medications would be critical to effective delivery of the drug with the limited window of absorption and many patients may thus fail to take their medication correctly resulting in ineffective treatment of their diabetes.

Prior Art Gastro-Retentive Systems

It would be desirable to provide a dosage form that inherently has the property of extended gastric residence, possessing some resistance to the pattern of waves of

motility present in the gastrointestinal tract that serve to propel material through it. There have been many attempts to provide for this, with varying degrees of success.

5

Possible approaches described in prior art include:

(1) Floating or buoyant systems:

These are designed to have a low density and thus
10 should float on gastric contents after
administration until the system either disintegrates
(and presumably the resultant particles empty from
the stomach) or the device absorbs fluid to the
point where its density is such that it loses
15 buoyancy and can pass more easily from the stomach
with a wave of motility responsible for gastric
emptying.

(2) Bioadhesive systems:

These are designed to imbibe fluid following
20 administration such that the outer layer becomes a
viscous, tacky material that adheres to the gastric
mucosa/mucus layer. This should encourage gastric
retention until the adhesive forces are weakened for
25 example by continuing hydration of the outer layer
of the device or by the persistent application of
shear.

(3) Swelling and expanding systems:

These are designed to be sufficiently small on
30 administration so as not to make ingestion of the
dosage form difficult (for example, less than
approximately 23 mm long and less than 11 mm wide
for an oval or capsule-shaped tablet). On ingestion
35 they rapidly swell or unfold to a size that
precludes passage through the pylorus until after
drug release has progressed to a required degree.

Gradual erosion of the system or its breakdown into smaller particles enables it to leave the stomach.

Re: (1) Buoyant/floating systems

5 Buoyant systems designed to float on the gastric contents have been designed where buoyancy is created by low density of the formulation components. For example, Watanabe et al (3) used low density shells such as spherical polystyrene foam particles in which polymer and
10 drug layers were loaded. Such a system has the required low density and will not need to disintegrate into small pieces to empty from the stomach, but may not have a controlled loss of density alternatively required for it to eventually exit from the stomach. It also has limited
15 capacity for loading with drug in the thin layers that can be applied around the polystyrene shells. It would be difficult to also layer large amounts of polymer on such a system to retard the release of very water soluble drugs.

Sheth described hydrodynamically balanced systems
20 including both capsules and tablets (4,5,6) which had low density to enable floating on the stomach contents and which slowly eroded after administration, losing buoyancy and being expelled from the stomach.

Buoyancy can also be combined with control of drug
25 release at different pH values to make for a device with better control in case of drugs with very marked dependency of solubility on pH (7); hence dissolution of contained drug depending on environment pH.

These approaches may be applicable to many drugs
30 dosed in doses of up to a maximum of a few hundred milligrams per day but may not be applicable to similar or higher dose levels of highly water soluble drugs. Where large amounts of polymer are needed to retard drug release as in the case of use of high water soluble drugs a capsule
35 dosage form may not be possible on grounds of size. Furthermore, the relatively homogenous distribution of drug in the tablet versions of this technology would not readily

control the burst effect seen with a very water soluble drug.

A bilayer tablet approach (8) where the buoyancy generation comes from a separate layer to the drug containing layer having a release rate controlling property might overcome some of the problems seen with the hydrodynamically balanced systems, but this type of system would probably only be able to carry low drug payloads due to size constraints.

Approaches involving in situ gas generation within the system, where the gas is trapped within the dosage form on generation, encouraging buoyancy, might offer improved control over degree, onset time and persistence of buoyancy. Ichikawa (9) described such a device with a drug loaded core surrounded by the gas generating layer, which in turn was surrounded by a polymeric layer responsible for controlling drug release from the system.

Such floating or buoyant dosage forms seem to have met with limited clinical success due to the requirement that such dosage forms be taken with a suitable amount of fluid (normal gastric contents could be as little as a few tens of milliliters so that the total amount of fluid thus available would not be conducive to performance of such systems even when taken with a draught of water). Davis et al (10) found no benefit of floating formulations over non-floating formulations when studied in vivo. Their performance may also be posture dependent. A patient sitting upright may ensure prolonged gastric residence of a buoyant dosage form, whereas a supine patient might allow ready presentation of the floating dosage form to the pylorus and thus allow rapid exit of the dosage form from the stomach (11). The physical size of such dosage forms seems to be as important if not more important as ability to float in encouraging prolonged gastric residence. Hence, floating/buoyant dosage forms might be expected to only have limited applications.

Re: (2) Bioadhesive systems

Polycarbophil has been identified as a suitable polymer for encouraging adhesion of orally administered dosage forms to the gastric mucosa, thereby prolonging residence time for a system designed to slowly deliver drug to absorptive sites in the proximal small intestine (Longer et al, J. Pharm. Sci., 74, 406-411 (1985)). The success seen in animal models with such systems has been found not to translate to human subjects due to differences in mucous amounts, consistency and turnover between animals and humans. Bioadhesive systems allow dosage forms to adhere to mucous, not mucosa. The mucous layer in humans would appear to slough off readily, carrying any dosage form with it. Therefore, bioadhesive dosage forms would not appear to offer a solution for extended delivery of drug over a period of more than a few hours to the upper small intestine in humans.

Re: (3) Swelling/expanding systems

Other solutions to encouraging prolonged gastric residence have included dosage forms that unfold rapidly within the stomach to a size that resists gastric emptying. Such systems retain their integrity for an extended period and will not empty from the stomach at all until breakdown into small pieces occurs. Caldwell (12) describes a cross shaped device made of erodible polymer and loaded with drug which is folded and inserted into a hard gelatin capsule. Following oral administration the gelatin shell disintegrates and the folded device opens out. With a minimum size of 1.6cm and a maximum size of 5cm it will not pass from the stomach through the pylorus until the polymer erodes to the point where the system is sufficiently small that it can be passed from the stomach. Such a system may in fact obstruct the pylorus or even open earlier or later than intended possibly causing obstruction in the esophagus or small intestine. As such, it may represent a potential hazard to the patient.

An alternate approach to using size to modulate gastric residence of a dosage form is to use a hydrophilic erodible polymer system that is of a convenient size for administration to humans. On imbibing fluid the system
5 swells over a short period of time to a size that will encourage prolonged gastric retention, allowing sustained delivery of contained drug to absorption sites in the upper gastrointestinal tract. Because these systems are made of an erodible and hydrophilic polymer or polymer mixture,
10 they readily erode over a reasonable time period to pass from the stomach. The time period of expansion is such that this will not occur in the esophagus and if the system passes into the intestine in a partially swollen state, the erodibility and elastic nature of the hydrated polymer will
15 eliminate the chance of intestinal obstruction by the device.

Mamajek et al, U.S. Patent No. 4,207,890, describes a system wherein a drug release rate controlling (metering) component and a swelling component are mixed with drug
20 enclosed within a membrane. The swelling component draws in fluid through the membrane, which maintains system integrity during its functioning, and the drug metering component controls the rate of drug release through the membrane.

25 Urquart (13) describes a different approach which consists of a matrix of hydrogel that imbibes fluid to swell the system so that it reaches a size encouraging prolonged gastric retention. This matrix surrounds a plurality of tiny pills consisting of drug with a release
30 rate controlling wall of fatty acid and wax surrounding each of the pills.

Shell (14,15,16) has described systems for delivering drugs for the treatment of diseases of the upper gastrointestinal tract or for delivering drugs that might
35 be irritating or injurious to the gastrointestinal mucosa. A swelling hydrogel polymer has embedded within it drug particles that dissolve once the hydrogel matrix is

hydrated. The swollen matrix is of a size to encourage gastric retention but only dissolved drug reaches the mucosa and this can be delivered in a sustained manner. Such a system thus does not insult the mucosa with solid
5 particles of irritant drug and is suitable for delivering drug to upper gastrointestinal tract. These systems only apply in case of drugs of limited water solubility.

In the case of metformin, it is desirable to provide a dosage form that allows extended delivery of the drug and
10 has a prolonged gastric residence via swelling of the system rather than unfolding or expanding of a folded device, and that may be manufactured on a commercial scale. The prolonged gastric residence time is required due to the window of absorption seen with metformin.

15 Another problem for extended delivery of metformin is its very high water solubility. High levels of polymer would be needed if many prior art approaches to provide the required release rate are employed. This could result in a rapid and variable initial release (burst) of drug from an
20 extended release dosage form. The latter will thus give rise to difficulty in providing a true control of drug release and minimal inter-patient variability in drug plasma levels (arising from the possibility of variable burst of drug from tablets given to different patients).

25

Prior Art Controlled Release Systems for Very Soluble Drugs

Typical prior art techniques for creating a controlled release oral dosage form would involve either
30 matrix systems or multi particulate systems. Matrix systems may be formulated by homogeneously mixing drug with hydrophilic polymers, such as hydroxypropylmethylcellulose, hydroxypropylcellulose, polyethylene oxide, carbomer, certain methacrylic acid derived polymers, sodium alginate,
35 or mixtures of components selected from these, and compressing the resultant mixture into tablets (employing some other excipients where needed). Hydrophobic polymers,

such as ethyl cellulose, certain polymeric methacrylic acid esters, cellulose acetate butyrate, poly(ethylene-co-vinyl-acetate) may be uniformly incorporated with the above materials to give additional control of release. A further
5 alternative involves embedding drug within a wax based tablet, by granulation or simply mixing of drug with a wax, such as carnauba wax, microcrystalline wax or commercially available purified fatty acid esters. As noted above, it may not be possible to use these approaches with very
10 highly water soluble drugs.

Multi particulate systems consist of a dosage form based on a plurality of drug loaded spheres, prepared by layering drug onto a core, usually a sugar-starch mixture sphere of around 0.8mm diameter, until a sufficient level
15 is reached, and then providing a drug release barrier around the drug-loaded sphere. Drug-loaded spheres can also be made by wet massing a mixture of drug and excipients, forcing the wet mass through a perforated screen to form short strands which are rounded in a
20 spheronisation apparatus before drying and having the drug release barrier applied. The drug release barrier can be a wax, such as carnauba wax or glyceryl fatty acid esters, or a polymeric barrier, such as a mixture of ethyl cellulose and hydroxypropylmethylcellulose. These work well for
25 moderately soluble drugs with doses in the units of milligrams to less than a few hundred milligrams per day.

In several examples, prior art systems seem to provide a controlled release formulation of a very water soluble drug by improving the multi particulate system
30 approach. Fisher discloses a multi particulate system for highly soluble drugs especially opiate agonists (17) based on drug containing cores surrounded by a drug release controlling barrier which has the property of being partially soluble at a highly acidic pH.

35 Hansraj (18) coats drug loaded cores with methacrylic or acrylic acid derived polymers whose properties are modified by inclusion of at least one

anionic surfactant. In such a system, drug release of highly water soluble drugs is controlled without having to resort to the use of thick coatings on the release rate controlling layer.

5 Rollet (19) achieves prolonged release of a drug from a multi particulate formulation based on fine particles of hydrophilic and hydrophobic silicas or silicates. Presumably, this system would function for drugs of high water solubility.

10 Multi particulate systems are usually filled into capsules to provide unit dose forms because of the damage caused to such particles in trying to compress them into tablets. Total dose contained in a single unit is constrained by the loading possible in a hard gelatin
15 capsule of easily swallowable size and is usually not more than a few hundred milligrams.

 Single unit controlled release systems applicable to highly water soluble drugs include the application of multiple layers around a dose form as described by Howard
20 (20). Where coating is not employed, special mixtures of polymers or formation of a complex with the drug have been used. Macrae (21) uses mixtures of polyethylene oxide and hydroxypropylmethylcellulose with optional enteric polymers to produce a constant release rate for highly water soluble
25 drugs. Belenduik (22) combines the highly water soluble drug with a hydrophilic polymer based on acrylic acid and disperses this in a hydrophobic matrix.

 Variations of Alza osmotic systems have been described suitable for highly water soluble drugs such as
30 venlafaxine hydrochloride (23). These systems need two layers, a drug layer and an osmotically driven displacement layer all surrounded by a water permeable/drug impermeable membrane with an exit passage in this membrane for the drug.

35 Granules of highly water soluble clavulanate were prepared (24) having to employ a barrier layer of a hydrophobic waxy material in order to provide for

controlled release of this material when co-formulated with controlled release amoxycillin trihydrate granules in capsule or compressed tablet.

5

Description of the Invention

In accordance with the present invention, a novel way has been found of formulating drug with high water solubility and a limited window of absorption such as metformin or a salt thereof which has a window of
10 absorption in the upper gastrointestinal tract, to provide a dosage form that inherently has prolonged gastric residence. This is accomplished (a) without need for co-administration of a drug such as propantheline, and (b) without need for low density formulation or gas generation
15 within the formulation. The formulation of the invention (a) achieves extended gastric residence by virtue of size but will degrade in vivo so as not to have potential for causing gastric or intestinal obstruction, and (b) controls drug release adequately where the initial burst of drug is
20 under control. The formulations of the invention will provide for an extended release formulation of drug with minimal interpatient variability in pharmacokinetic parameters.

The invention is applicable to all drugs having high
25 water solubility and a limited window of absorption.

The biphasic controlled release delivery system of the invention is a heterogeneous two phase system which includes (1) an inner solid particulate phase in the form of individual granules or particles containing (a) drug
30 which has a high water solubility and a limited window of absorption (such as in the upper gastrointestinal tract), and (b) an extended release material formed of one or more hydrophilic polymers, and/or one or more hydrophobic polymers, and/or one or more other type hydrophobic
35 compounds (such as one or more waxes, fatty alcohols and/or fatty acid esters), and (2) an outer solid continuous phase in which granules or particles of inner solid particulate

phase are dispersed and embedded, the outer solid continuous phase including an extended release material formed of one or more hydrophilic polymers, and/or one or more hydrophobic polymers, and/or one or more other type
5 hydrophobic compounds (such as one or more waxes, fatty alcohols and/or fatty acid esters).

The biphasic controlled release formulation of the invention is particularly adapted for delivery of high water soluble drugs, such as metformin and pharmaceutically
10 acceptable salts thereof, in controlled and extended manner without significant initial burst of drug, and wherein release of drug (liberated from the individual dispersed particles forming the inner solid particulate phase) is effectively controlled. Drug upon being released from the
15 particles of the inner phase, in effect, migrates through the outer solid continuous phase and then is released from the formulation into the upper gastrointestinal tract to be available for absorption.

As indicated, the inner solid particulate phase will
20 be formed of individual discrete particles or granules each of which contains drug and one or more polymeric materials and/or other hydrophobic-type materials. In effect, the components of the inner solid particulate phase are in particulate association without having a barrier layer
25 around the individual particles or granules.

The outer solid continuous phase is preferably a continuous phase or matrix having the particles or granules including drug (forming the inner solid phase) dispersed throughout and embedded in the continuous outer solid
30 phase.

In addition, in accordance with the present invention, a method for treating diabetes is provided wherein the biphasic controlled release formulation of the invention containing an antidiabetic pharmaceutical is
35 administered to a patient in need of treatment. The antidiabetic pharmaceutical employed is a biguanide,

preferably metformin or a pharmaceutically acceptable salt thereof such as the hydrochloride, fumarate of succinate.

The term "extended release material" as present in the inner solid particulate phase and the outer solid continuous phase refers to one or more hydrophilic polymers and/or one or more hydrophobic polymers and/or one or more other type hydrophobic materials, such as, for example, one or more waxes, fatty alcohols and/or fatty acid esters. The "extended release material" present in the inner solid particulate phase may be the same as or different from the "extended release material" present in the outer solid continuous phase. However, it is preferred that the "extended release material" present in the inner solid particulate phase be different from the "extended release material" present in the outer solid continuous phase.

The term "high water solubility" or similar term when characterizing a drug, medicament or pharmaceutical for use in the formulation of the invention refers to a solubility in water at ambient temperature of at least about 50 mg/ml H₂O, preferably at least about 100 mg/ml H₂O or more, and more preferably greater than 150 mg/ml.

The term "limited window of absorption" or similar term when characterizing a drug, medicament or pharmaceutical for use in the formulation of the invention refers to an oral bioavailability of less than about 75%, usually less than about 60%, usually decreasing with increasing dose, and almost invariably having permeability/transit time limited absorption.

The biphasic controlled release system of the invention will include the inner solid particulate phase in a weight ratio to the outer solid continuous phase within the range from about 0.5:1 to about 4:1, preferably from about 0.8:1 to about 2:1.

The inner solid particulate phase will contain drug in an amount within the range from about 10 to about 98% by weight, preferably from about 15 to about 95% by weight, and hydrophilic polymers and/or hydrophobic polymers and/or

other hydrophobic material in an amount within the range from about 5 to about 95% by weight, preferably from about 7 to about 85% by weight, the above % being based on the weight of the inner solid particulate phase. Where
5 mixtures are employed, the hydrophilic polymer will be employed in a weight ratio to hydrophobic polymer and/or other hydrophobic material within the range from about 0.05:1 to about 19:1, preferably from about 0.1:1 to about 10:1.

10 The particles or granules of the inner solid particulate phase will have a mean particle size within the range from about 30 μm to about 0.8 mm, and preferably from about 50 μm to about 0.5 mm.

The outer solid continuous phase may contain
15 mixtures of hydrophilic polymer and hydrophobic polymer and/or other hydrophobic material in a weight ratio of hydrophilic polymer to hydrophobic polymer or other hydrophobic material within the range from about 200:1 to about 0.05:1, preferably from about 100:1 to about 0.1:1.

20 Hydrophilic polymers which may be employed in the inner solid particulate phase and/or outer solid continuous phase include, but are not limited to hydroxypropylmethylcellulose, hydroxypropylcellulose, sodium carboxymethylcellulose, carboxymethylcellulose
25 calcium, ammonium alginate, sodium alginate, potassium alginate, calcium alginate, propylene glycol alginate, alginic acid, polyvinyl alcohol, povidone, carbomer, potassium pectate, potassium pectinate, and the like.

Hydrophobic polymers which may be employed in the
30 inner solid particulate phase and/or outer solid continuous phase include, but are not limited to ethyl cellulose, hydroxyethylcellulose, ammonio methacrylate copolymer (Eudragit RL[™] or Eudragit RS[™]), methacrylic acid copolymers (Eudragit L[™] or Eudragit S[™]), methacrylic acid-
35 acrylic acid ethyl ester copolymer (Eudragit L 100-5[™]), methacrylic acid esters neutral copolymer (Eudragit NE

30D™), dimethylaminoethylmethacrylate-methacrylic acid esters copolymer (Eudragit E 100™), vinyl methyl ether/maleic anhydride copolymers, their salts and esters (Gantrez™).

5 Other hydrophobic materials which may be employed in the inner solid particulate phase and/or outer solid continuous phase include, but are not limited to waxes such as beeswax, carnauba wax, microcrystalline wax, and ozokerite; fatty alcohols such as cetostearyl alcohol, 10 stearyl alcohol; cetyl alcohol and myristyl alcohol; and fatty acid esters such as glyceryl monostearate, glycerol monooleate, acetylated monoglycerides, tristearin, tripalmitin, cetyl esters wax, glyceryl palmitostearate, glyceryl behenate, and hydrogenated castor oil.

15 Where hydrophilic polymers and/or hydrophobic polymers are used in the inner solid particulate phase and/or the outer solid continuous phase, such polymers can be ionic or non-ionic, preferably ionic for the inner solid particulate phase and preferably non-ionic for the outer 20 solid continuous phase.

 Preferred ionic polymers for use in the inner solid particulate phase include sodium alginate, carbomer (Carbopol™), calcium carboxymethylcellulose, or sodium 25 carboxymethylcellulose, xanthan gum, methacrylic acid-acrylic acid ethyl ester copolymer, dimethylaminoethylmethacrylate-methacrylic acid esters copolymer, cellulose acetate phthalate, hydroxypropyl-methylcellulose phthalate, hydroxypropylmethylcellulose trimellitate, and hydroxypropylmethylcellulose maleate, 30 with sodium carboxymethylcellulose being particularly preferred.

 Preferred non-ionic polymers for use in the outer solid continuous phase are those which assure rapid 35 hydration of the outer solid continuous phase to minimize a variable and significant burst of drug, yet effectively control the release of drug being liberated from the discrete particles or granules forming the inner solid

particulate phase. The liberated drug will migrate through the non-ionic polymers forming the outer solid continuous phase before being released from the dosage form and being available for absorption. Preferred polymers for the outer solid phase with the appropriate hydration characteristics include hydroxypropylmethyl cellulose 2910 USP (hydroxypropylmethylcellulose with a methoxyl content of 19-24% and a hydroxypropyl content of 7-12%), viscosity grades ranging from about 4000 to about 100,000 cps and hydroxypropylmethylcellulose 2208 USP (hydroxypropylmethylcellulose with a methoxyl content of 28-30% and a hydroxypropyl content of 7-12%), viscosity grades ranging from about 3 to about 150 cps, and microcrystalline cellulose. In particular preferred embodiments of the outer solid phase, the above preferred polymers are used in admixture in weight ratios of hydroxypropylmethylcellulose 2910 USP:hydroxypropylmethylcellulose 2208 USP within the range from about 25:1 to about 50:1, preferably from about 30:1 to about 40:1.

Preferred biphasic controlled extended release delivery systems in accordance with the present invention are as follows.

	% by Weight of Inner_ Solid Particulate Phase
A. <u>Inner Solid Particulate Phase</u> 25 (1) Metformin HCl (or other salt such as succinate or fumarate)	55 to 98
30 (2) Polymer or Hydrophobic material: ethylcellulose or sodium carboxymethylcellulose or glyceryl monostearate (Average Particle Size of granules forming inner solid particulate 35 phase: 0.05 to 2.0 mm)	5 to 45

		% by Weight of Outer_ Solid Continuous Phase
B.	<u>Outer Solid Continuous Phase</u>	
	(1) Hydroxypropylmethyl- cellulose 2208USP	60 to 100
5	(100,000 cps)	
	(2) Microcrystalline cellulose	0 to 30
	(3) Hydroxypropylmethyl cellulose 2910 USP (5 cps)	1 to 30
	Weight Ratio of Inner Solid Phase:	0.5:1 to 1.5:1
10	Outer Solid Phase	
		% by Weight of Outer_ Solid Continuous Phase
C.	<u>Optional Ingredients</u>	
	Lubricant - Mg Stearate	0.02 to 1
15		

The preferred drug (having high water solubility) for use herein is metformin or pharmaceutically acceptable salts thereof, including the hydrochloride salt and dibasic salts such as metformin (2:1) fumarate and metformin (2:1) succinate as described in pending U.S. Application Serial No. (not yet available) filed December, 1997 (internal code LA19) which is incorporated herein by reference. Other biguanides may be used such as phenformin or buformin or pharmaceutically acceptable salts thereof.

Most preferred are the metformin hydrochloride salt, metformin (2:1) succinate salt, and metformin (2:1) fumarate salt.

Where desired, metformin or a salt thereof may be used in combination with another antihyperglycemic agent which may be administered orally in the same dosage form in accordance with the invention, a separate oral dosage form or by injection.

It is believed that the use of the metformin or salt thereof in combination with another anti-hyperglycemic agent produces antihyperglycemic results greater than that possible from each of these medicaments alone and greater

than the combined additive anti-hyperglycemic effects produced by these medicaments.

The other antihyperglycemic agent may be an oral antihyperglycemic agent preferably a sulfonyl urea such as glyburide (also known as glibenclamide), glimepiride (disclosed in U.S. Patent No. 4,379,785), glipizide, gliclazide or chlorpropamide, other known sulfonylureas or other antihyperglycemic agents which act on the ATP-dependent channel of the β -cells, with glyburide being preferred.

The metformin or salt thereof will be employed in a weight ratio to the sulfonyl urea in the range from about 300:1 to about 50:1, preferably from about 250:1 to about 75:1.

The oral antihyperglycemic agent may also be a glucosidase inhibitor such as acarbose (disclosed in U.S. Patent No. 4,904,769) or miglitol (disclosed in U.S. Patent No. 4,639,436), which may be administered in a separate oral dosage form.

The metformin salt of the invention will be employed in a weight ratio to the glucosidase inhibitor within the range from about 300:1 to about 2:1, preferably from about 200:1 to about 25:1.

The metformin or salt thereof may be employed in combination with a thiazolidinedione oral anti-diabetic agent (which has an insulin sensitivity effect in NIDDM patients) such as troglitazone (Warner-Lambert's Rezulin®, disclosed in U.S. Patent No. 4,572,912), zorglitazone (SKB), pioglitazone (Takeda), Mitsubishi's MCC-555 (disclosed in U.S. Patent No. 5,594,016) Glaxo-Welcome's GL-262570, englitazone (CP-68722, Pfizer) or darglitazone (CP-86325, Pfizer).

The metformin or salt thereof will be employed in a weight ratio to the thiazolidinedione in an amount within the range from about 75:1 to about 0.1:1, preferably from about 5:1 to about 0.5:1.

The sulfonyl urea and thiazolidinedione in amounts of less than about 150 mg oral anti-diabetic agent may be incorporated in a single tablet with the biphasic controlled release formulation of the invention as a
5 separate rapidly dissolving layer.

The metformin or salt thereof may also be employed in combination with a non-oral antihyperglycemic agent such as insulin or with glucagon-like peptide-1 (GLP-1) such as GLP-1(1-36) amide, GLP-1(7-36) amide, GLP-1(7-37) (as
10 disclosed in U.S. Patent No. 5,614,492 to Habener, the disclosure of which is incorporated herein by reference), which may be administered via injection, or by transdermal or buccal devices.

In addition, in accordance with the present
15 invention a method is provided for treating hyperglycemia including Type II diabetes (NIDDM) and/or Type I diabetes (IDDM) wherein a therapeutically effective amount of the biphasic formulatin of the invention containing metformin or a salt thereof, optionally in combination with another
20 antihyperglycemic agent, is administered to a patient in need of treatment.

Where present, the sulfonyl ureas, such as glyburide, glimepiride, glipyrider, glipizide, glipizide, chlorpropamide and gliclazide and the glucosidase
25 inhibitors acarbose or miglitol may be employed in formulations as described above and in amounts and dosing as indicated in the Physician's Desk Reference.

Where present, the thiazolidinedione anti-diabetic agent may be employed in amounts within the range from
30 about 0.01 to about 2000 mg/day which may be administered in single or divided doses one to four times per day.

Where present insulin may be employed in formulations, amounts and dosing as indicated by the Physician's Desk Reference.

35 Where present GLP-1 peptides may be administered in oral buccal formulations, by nasal administration or parenterally as described in U.S. Patent Nos. 5,346,701

(TheraTech), 5,614,492 and 5,631,224 which are incorporated herein by reference.

The following additional type high water soluble drugs may be employed in the biphasic controlled release
5 delivery system of the invention:

antihypertensives and antidepressants related to guanethidine (as disclosed in U.S. Patent No. 2,928,829) and related to guanoxyfen (as disclosed in BE612362);

antibiotics and viricides such as related to
10 amidinomycin (as disclosed in JP 21,418);

stallimycin (as disclosed in DE 1,039,198);

Arphamenine B (as disclosed in published European Patent Application 85/133550A2);

chitinovorin-A (as disclosed in published European
15 Patent Application 85/150,378A2 and U.S. Patent No.,
4,723,004);

streptomycin (as disclosed in U.S. Patent No.
2,868,779);

SB-59 (as disclosed in Justus Liebigs, Ann. Chem.
20 (1973) 7, 1112-1140);

TAN-1057-A (as disclosed in U.S. Patent No.
4,971,965);

streptoniazid (as disclosed in J. Am. Chem. Soc.
(1953) 75, 2261);

25 immunostimulants related to

ST-789 (as disclosed in published European Patent Application 88/260588);

peptide hydrolase inhibitors related to

nafamastat (as disclosed in U.S. Patent No.
30 4,454,338);

gabexate (as disclosed in U.S. Patent No.
3,751,447);

sepimostat (as disclosed in U.S. Patent Nos.
4,777,182 and 4,820,730);

35 Factor Xa inhibitors related to

DX-9065a (as disclosed in published European Patent Application 92/0540051);

anti-inflammatory agents related to paranyline as disclosed in U.S. Patent No. 2,877,269;

peptidyl aldehydes (as disclosed in WO94/13693);

antianaphylactics related to GMCHA-TBP (Batebulast)

5 (as disclosed in U.S. Patent No. 4,465,851);

anti-ulcer agents related to

benexate (as disclosed in U.S. Patent No.

4,348,410);

deoxyspergualin (as disclosed in U.S. Patent Nos.

10 4,518,532, 4,658,058 and 4,983,328); and arginine.

Other water-soluble drugs suitable for use herein include peptides preferably have a molecular weight from about 100 to 10,000, more preferably from about 100 to about 6,000 and having from 2 to 35 amino acid moieties.

15 Higher molecular weight peptides, even those with a molecular weight of above 10,000, up to about 50,000, may also be accommodated in biphasic formulations of the present invention.

Suitable small peptides have from about 2 to about
20 10, more preferably from about 2 to about 6 amino acid moieties. Preferred small peptides include the fibrinogen receptor antagonists (RGD containing peptides) which are tetrapeptides with an average molecular weight of about 600. These peptide antagonists are highly potent platelet
25 aggregation inhibitors at plasma levels as low as 1 pmol/mL. Preferred fibrinogen antagonists include the peptide cyclo(S,S)-N^a-acetyl-Cys-(N^a-methyl)Arg-Gly-Asp-Pen-NH₂ (Ali et al, EP 0341915, whose disclosure is herein incorporated by reference) and the peptide cyclo(S,S)-(2-
30 mercapto)benzoyl-(N^a-methyl)Arg-Gly-Asp-(2-mercapto)-phenylamide (EP 0423212, whose disclosure is herein incorporated by reference). Other fibrinogen antagonists useful in the present invention are those peptides disclosed by Pierschbacher et al, WO 89/05150
35 (US/88/04403); Marguerie, EP 0275748; Adams et al, U.S. 4,857,508; Zimmerman et al, U.S. 4,683,291; Nutt et al, EP 0410537, EP 0410539, EP 0410540, EP 0410541, EP 0410767, EP

0410833, EP 0422937 and EP 0422938; Ali et al, EP 0372486; Ohba et al, WO 90/02751 (PCT/JP89/00926); Klein et al, U.S. 4,952,562; Scarborough et al, WO 90/15620 (PCT/US90/03417); Ali et al, PCT/US90/06514 and PCT/US92/00999; the peptide-
5 like compounds disclosed by Ali et al, EP 0381033 and EP 0384362; and the RGD peptide cyclo-N^a-acetyl-Cys-Asn-Dtc-Amf-Gly-Asp-Cys-OH (in which Dtc is 4,4'-dimethylthiazolidine-5-carboxylic acid and Amf is 4-aminomethylphenylalanine).

10 The RGD peptide may be usefully included in the formulation of the invention in an amount up to about 600 mg/g of the hydrophilic phase or from 0.1 to 60 mg/g of the formulation.

Other peptides useful in the present invention
15 include, but are not limited to, other RGD containing peptides such as those disclosed by Momany, U.S. 4,411,890 and U.S. 4,410,513; Bowers et al, U.S. 4,880,778, U.S. 4,880,777, U.S. 4,839,344; and WO 89/10933 (PCT/US89/01829); the peptide Ala-His-D-Nal-Ala-Trp-D-Phe-
20 Lys-NH₂ (in which Nal represents β -naphthylalanine) and the peptides disclosed by Momany, U.S. 4,228,158, U.S. 4,228,157, U.S. 4,228,156, U.S. 4,228,155, U.S. 4,226,857, U.S. 4,224,316, U.S. 4,223,021, U.S. 4,223,020, U.S. 4,223,019 and U.S. 4,410,512.

25 Other suitable peptides include hexapeptides such as the growth hormone releasing peptide (GHRP) His-D-Trp-Ala-Trp-D-Phe-Lys-NH₂, (Momany, U.S. 4,411,890, the disclosure of which is herein incorporated by reference in its entirety). This may usefully be included in an amount up
30 to about 250 mg/g of the hydrophilic phase or from 0.1 to 25 mg/kg of the formulation.

Suitable larger polypeptides and proteins for use in the controlled release formulations of the present invention include insulin, calcitonin, elcatonin,
35 calcitonine gene related peptide and porcine somatostatin as well as analogs and homologs thereof. Other suitable larger polypeptides include those disclosed by

Pierschbacher et al, U.S. 4,589,881 (>30 residues); Bittle et al, U.S. 4,544,500 (20-30 residues); and Dimarchi et al, EP 0204480 (>34 residues).

Other type of compounds useful in the present invention include analogs or homologs of LHRH which display potent LH releasing activity or inhibit the activity of LHRH; analogs or homologs of HP5 which possesses hematopoietic activity; analogs or homologs of endothelin which possess hypotensive activity; analogs or homologs of enkephalin which have antinociceptive activity; analogs or homologs of chloreocystokinin; analogs or homologs of cyclosporin A which have immunosuppressive activity; analogs or homologs of atrial natriuretic factor; peptidergic antineoplastic agents; analogs or homologs of gastrin releasing peptide; analogs or homologs of somatostatin; gastrin antagonists; bradykinin antagonists; neurotensin antagonists; bombesin antagonists; oxytocin agonists and antagonists; vasopressin agonists and antagonists; hirudin analogs and homologs; analogs and homologs of the cytoprotective peptidecyclolinopeptide; alpha MSH analogs; analogs, and homologs of MSH releasing factor (Pro-Leu-Gly-NH₂); peptides which inhibit collagenase; peptides which inhibit elastase, peptides which inhibit renin; peptides which inhibit HIV protease; peptides which inhibit angiotensin converting enzyme; peptides which inhibit chymases and tryptases and peptides which inhibit blood coagulation enzymes.

Other suitable drugs include non-peptide therapeutic agents such as antibiotics, antimicrobial agents, antineoplastic agents, cardiovascular and renal agents, anti-inflammatory, immunosuppressive and immunostimulatory agents and CNS agents.

Preferably, the water-soluble drug is metformin or salt thereof as described above.

The biphasic controlled release formulation of the present invention can be administered to various mammalian

species, such as dogs, cats, humans, etc., in need of such treatment.

The biphasic controlled release system of the invention can be incorporated in a conventional systemic dosage form, such as a tablet or capsule. The above dosage forms may also include the necessary physiologically acceptable carrier material, excipient, lubricant, buffer, antibacterial, bulking agent (such as mannitol), anti-oxidants (ascorbic acid or sodium bisulfite) or the like.

The dose administered must be carefully adjusted according to the age, weight, and condition of the patient, as well as the route of administration, dosage form and regimen, and the desired result. In general, the dosage forms of formulation containing metformin or salt thereof (whether by itself or with another antihyperglycemic agent) described above may be administered in amounts as described for metformin hydrochloride (Bristol-Myers Squibb Company's Glucophage®) as set out in the Physician's Desk Reference.

The combination of the metformin or salt thereof and the other antihyperglycemic agent may be formulated separately or, where possible, in a single formulation employing conventional formulation procedures.

The various formulations of the invention may optionally include one or more fillers or excipients in an amount within the range of from about 0 to about 90% by weight and preferably from about 1 to about 80% by weight such as lactose, sugar, corn starch, modified corn starch, mannitol, sorbitol, inorganic salts such as calcium carbonate and/or cellulose derivatives such as wood cellulose and microcrystalline cellulose.

One or more binders may be present in addition to or in lieu of the fillers in an amount within the range of from about 0 to about 35% and preferably from about 0.5 to about 30% by weight of the composition. Examples of such binders which are suitable for use herein include polyvinylpyrrolidone (molecular weight ranging from about 5000 to about 80,000 and preferably about 40,000), lactose,

starches such as corn starch, modified corn starch, sugars, gum acacia and the like as well as a wax binder in finely powdered form (less than 500 microns) such as carnauba wax, paraffin, spermaceti, polyethylenes or microcrystalline wax.

Where the composition is to be in the form of a tablet, it will include one or more tableting lubricants in an amount within the range of from about 0.2 to about 8% and preferably from about 0.5 to about 2% by weight of the composition, such as magnesium stearate, stearic acid, palmitic acid, calcium stearate, talc, carnauba wax and the like. Other conventional ingredients which may optionally be present include preservatives, stabilizers, anti-adherents or silica flow conditioners or glidants, such as Syloid brand silicon dioxide as well as FD&C colors.

Tablets of the invention may also include a coating layer which may comprise from 0 to about 15% by weight of the tablet composition. The coating layer which is applied over the outer solid phase containing particles of inner solid phase embedded therein may comprise any conventional coating formulations and will include one or more film-formers or binders, such as a hydrophilic polymer like hydroxypropylmethylcellulose, and/or a hydrophobic polymer like methacrylic acid esters neutral polymer, ethyl cellulose, cellulose acetate, polyvinyl alcohol-maleic anhydride copolymers, β -pinene polymers, glyceryl esters of wood resins and the like and one or more plasticizers, such as triethyl citrate, diethyl phthalate, propylene glycol, glycerin, butyl phthalate, castor oil and the like. Both core tablets as well as coating formulations may contain aluminum lakes to provide color.

The film formers are applied from a solvent system containing one or more solvents including water, alcohols like methyl alcohol, ethyl alcohol or isopropyl alcohol, ketones like acetone, or ethylmethyl ketone, chlorinated hydrocarbons like methylene chloride, dichloroethane, and 1,1,1-trichloroethane.

Where a color is employed, the color will be applied together with the film former, plasticizer and solvent compositions.

It will be recognized by one of skill in the art
5 that the amount of drug required for therapeutic effect on administration will, of course, vary with the agent chosen, the nature and severity of the condition and the animal undergoing treatment, and is ultimately at the discretion of the physician. Furthermore, the optimal quantity and
10 spacing of individual dosages of a drug will be determined by the nature and extent of the condition being treated, the form, route and site of administration, the particular patient being treated and that such optima can be determined by conventional techniques. It will also be
15 appreciated that the optimal course of treatment, this is, the number of doses given, can be ascertained by those skilled in the art using conventional course of treatment determination tests.

As indicated, the preferred highly water-soluble
20 drug will be metformin or a salt thereof, which will be employed in a dosage range from about 2 to about 43 mg/kg, preferably about 3 to about 36 mg/kg and more preferably from about 4.5 to about 30 mg/kg (or from about 150 to about 3000 mg, preferably from about 250 to about 2500 mg)
25 on a regimen in single or 2 to 4 divided daily doses.

The biphasic controlled release formulation of the invention may be prepared in accordance with the following method of the invention.

A mixture of medicament (preferably metformin HCl)
30 and hydrophilic polymer and/or hydrophobic polymer and/or other hydrophobic material are dispersed/dissolved in a suitable solvent such as water or an inert organic solvent such as ethanol, isopropanol, acetone or dichloromethane or appropriate mixtures of two or more thereof, to produce a
35 substantially uniform granulation. The granulation is dried and passed through a 0.5 to 2 mm aperture screen to break down agglomerates.

The resulting dry granules are blended with hydrophilic polymer and/or hydrophobic polymer and/or other hydrophobic material. The resulting mix usually with lubricant is pressed into tablets or filled into capsules.

5 The finished dosage form is either a compressed tablet or a hard gelatin capsule, preferably a tablet. The tablet may be optionally film coated. The total amount of drug per dosage unit would be such as to offer a dosage form of convenient size for patients, but following
10 ingestion would remain (or swell to, by hydration of the polymers used in the fabrication of the tablet) a size that does not easily pass through the pylorus (15mm or greater) when taken with a meal. As the tablet swells up to approximately three times its dry size following hydration,
15 drug loads of up to 750 mg are possible, dependent upon the actual characteristics of the individual drug. Gradual erosion of the polymers of the formulation over a period of up to 15 hours ensures that the dosage form does not produce a gastrointestinal obstruction.

20 Useful metformin formulations of the invention show the following drug release characteristic when tested in vitro.

	<u>Time (hours)</u>	<u>% released</u>
25	1	28-39
	2	43-57
	3	53-70
	5	70-88
	7	80-98
30	10	>85

The following Examples represent preferred embodiments of the invention.

Example 1Biphasic Metformin HCl Formulation

25g of ethylcellulose N10 NF was dissolved/
dispersed in 100 ml of 95% ethanol. This dispersion was
5 gradually added to 500 g of metformin hydrochloride in a
planetary mixer to produce a uniform damp granulation. The
granulation was dried at 55°C for one hour and passed
through a 0.8mm aperture screen to break down agglomerates.
The metformin-ethylcellulose granules (541g) were blended
10 with 351.5g of hydroxypropylmethylcellulose 2208 USP
(100,000 cps grade), 10g of hydroxypropylmethylcellulose
2910 USP (5 cps grade) and 100.5g of microcrystalline
cellulose in a planetary mixer for 10 minutes. Finally
this mix was lubricated with 1% w/w magnesium stearate and
15 compressed into capsule shaped tablets, each containing
500mg metformin hydrochloride. When subjected to in vitro
drug release testing, the following results were obtained.

	<u>Time (hours)</u>	<u>% metformin released</u>
20	1	38.1
	2	56.3
	3	69.5
	4	79.7
	5	87.4
25	6	93.1
	7	97.7
	8	100

Example 2Biphasic Metformin HCl Formulation

30 51g of sodium carboxymethylcellulose (Blanose 7HF)
was mixed with 500g of metformin hydrochloride and
granulated with 95% ethanol in a small planetary mixer.
The damp granulation was passed through a 2mm aperture
35 screen and then dried in an oven at 55°C for one hour. The
dried granulation (530g) was blended with 344g of
hydroxypropylmethylcellulose 2208 USP (100,000 cps grade),

9.5g of hydroxypropylmethylcellulose 2910 USP (5cps grade) and 100g of microcrystalline cellulose in a planetary mixer for 10 minutes. This blend was lubricated with 1% w/w magnesium stearate and compressed into capsule shaped tablets, each containing 500mg metformin hydrochloride. When the tablets were subjected to in vitro release testing the following results were obtained.

	<u>Time (hours)</u>	<u>% metformin released</u>
10	1	35.3
	2	51.4
	3	62.6
	4	70.7
	5	76.7
15	6	82.1
	7	85.3
	8	88.5
	10	92.6

20

Example 3Biphasic Metformin HCl Formulation

Metformin hydrochloride (502.5g) was mixed with sodium carboxymethylcellulose (Blanose 7HF) (50g) for five minutes in a small planetary mixer and sufficient purified water added with continued mixing to produce a damp granular mass. The wet granulation was dried at 60°C for 1 hour and then size reduced in a hammer mill. The granulation was dry blended with a mixture prepared from 385g of hydroxypropylmethylcellulose 2208 USP (100,000 cps grade), 10g of hydroxypropylmethylcellulose 2910 USP (5 cps grade) and 102g of microcrystalline cellulose in a planetary mixer for 10 minutes. Finally this mix was lubricated by mixing with 1% w/w magnesium stearate and compressed into capsule shaped tablets, each containing 0.5g of metformin hydrochloride. When tested for in vitro release of metformin the following results were obtained.

	<u>Time (hours)</u>	<u>% metformin released</u>
	1	33.1
	2	47.6
	3	57.5
5	4	65.1
	6	76.5
	8	84.3
	10	88.6

10

Example 4Biphasic Metformin HCl Formulation

200g of glycerol monostearate was heated to 70°C in a high shear mixer bowl and 199g of metformin hydrochloride was added and the mixer operated with impeller at 90 rpm and chopper at 215 rpm for 5 minutes. A further 796g of metformin hydrochloride was added gradually with continued mixing, maintaining the granulation at 70°C and with an increase in chopper speed first to 500 rpm for 13 minutes, then to 1000 rpm for a further 3 minutes. The bowl was then cooled to 60°C, the impeller speed was reduced to 20 rpm and the chopper speed increased to 2000 rpm. Cooling was continued with adjustment in impeller and chopper speed to eventually provide a cooled solid granulation. The cooled granulation was deagglomerated by passing through a 0.8mm screen.

540.5g of the granulation was blended with 350g of hydroxypropylmethylcellulose 2208 USP (100,000 cps grade), 10g hydroxypropylmethylcellulose 2910 USP (5 cps grade) and 100g of microcrystalline cellulose in a planetary mixer for 10 minutes. The blend was lubricated by blending with 1% w/w magnesium stearate and then compressed into capsule shaped tablets each containing 500mg metformin hydrochloride. When tested for in vitro release of metformin, the following results were obtained.

35

	<u>Time (hours)</u>	<u>% metformin released</u>
	1	32.4
	2	45.7
	3	55.8
5	4	63.7
	5	70.3
	6	75.7
	8	83.3
10	10	88.6

Example 5

Biphasic Metformin HCl Formulation

Tablets containing 500 mg metformin hydrochloride prepared according to Example 3 or Glucophage brand metformin hydrochloride 500 mg tablets was dosed (2 x 500 mg tablets) to 24 patients immediately after dinner. Blood samples were taken at 0, 0.5, 1, 2, 3, 4, 5, 6, 7, 8, 10, 12, 14, 16, 20, 24 hours and analyzed for metformin. The mean plasma profile demonstrated useful modification of drug release in vivo relative to the immediate release formulation and with no impact on bioavailability in contrast to other metformin extended release formulations reported in the literature.

Interpatient variability in pharmacokinetic parameters was acceptable as illustrated by the mean parameters (%CV) given in the table below:

Formulation	C _{max} (ng/ml)	AUC (inf) (ng.hr/ml)	T _{max} * (hr)	%UR
Glucophage	1226(16)	10128(14)	3.5(1,5)	43.3(20)
Example 3	978(13)	10483(21)	5(4,8)	42.7(18)

*median (min., max.)

The new formulations of the invention thus represent a useful advance in the administration of metformin hydrochloride to human in the treatment of diabetes.

Example 6Preparation of Metformin (2:1) Fumarate

Metformin base (8.71 moles) (prepared from the
5 hydrochloride salt via an ion-exchange column) was
dissolved in methanol/H₂O [5:1]. With stirring, a solution
of fumaric acid (4.05 moles) in ethanol was added over a
period of one hour under a nitrogen atmosphere at ambient
temperature (~20°C). Crystallization began to occur
10 immediately. After stirring the slurry for one hour at
ambient temperature, the product was filtered off, washed
with ethanol and dried under vacuum to afford the metformin
(2:1) fumarate salt as a free-flowing white crystalline
solid in 72 M% yield and melting point of 247-249°C.

15 The resulting metformin (2:1) fumarate salt had a
solubility in water (mg/ml) of 140, a hygroscopicity
measured at 95% relative humidity/25°C of less than 7%
moisture uptake at 6 hours, and a low compaction
susceptibility.

20 The so-formed metformin salt is used to prepare a
biphasic controlled release formulation employing the
procedure of Example 3.

Example 7Preparation of Metformin (2:1) Succinate

Metformin base (8.95 moles) (prepared from the
hydrochloride salt via an ion-exchange column) was
dissolved in methanol/H₂O [5:1]. With stirring, a solution
of succinic acid (4.42 moles) in ethanol was added over one
30 hour under a nitrogen atmosphere at ambient temperature
(~20°). Crystallization of the salt commenced shortly
after addition of the succinic acid solution. After
stirring the slurry for an hour at ambient temperature, the
product was filtered off, washed with ethanol and dried
35 under vacuum to form the metformin (2:1) succinate salt as
a free flowing white crystalline solid in 89 M% yield and
melting point of 246-247°C.

The resulting metformin (2:1) succinate salt had a solubility in water (mg/ml) of 95, a hygroscopicity measured at 95% relative humidity/25°C of less than 1% moisture uptake at 30 minutes, and a low compaction
5 susceptibility.

The so-formed metformin salt is used to prepare a biphasic controlled release formulation employing the procedure of Example 3.

10 REFERENCES

1. Vidon, N., Chaussade, S., Noel, M., Franchisseur, C., Huchet, B., Bernier, J.J. (1988), Diabetes Research and Clinical Practice, 4, 223-239.
2. Pentikainen, P.J. (1986), International Journal
15 of Clinical Pharmacology, Therapy and Toxicology, 24, 213-220.
3. Watanabe, S., Kayano, M., Ishino, Y., Miyao, K. (1976), U.S. Patent 3,976,764.
4. Sheth, P., Tossounian, J.L. (1979), U.S. Patent
20 4,140,755.
5. Sheth, P., Tossounian, J. L. (1978), U.S. Patent 4,126,672.
6. Sheth, P., Tossounian, J. L. (1979), U.S. Patent 4,167,558.
- 25 7. Dennis, A.B., Lee, K., Timmins, P. (1992), U.S. Patent 5,169,638.
8. Franz, M.R., Oth, M.P. (1993), U.S. Patent 5,232,704.
9. Ishikawa, M., Miyaka, Y., Watanabe, S. (1989),
30 U.S. Patent 4,844,905.
10. Davis, S.S., Stockwell, A.F., Taylor, J.J., Hardy, J.G., Whalley, D.R., Wilson, C.G., Bechgaard, H., Christensen, F.N. (1986) Pharm. Res., 3, 208-213.
11. Timmermans, J., Moes, A. (1994), J. Pharm.
35 Sci., 83, 18-24.
12. Caldwell, L.J., Gardener, C.R., Cargill, R.C. (1988), U.S. Patent 4,767,627.

13. Urquart, J., Theeuwes, F. (1984), U.S. Patent 4,434,153.
14. Shell, J.W. (1990), U.S. Patent 5,007,790.
15. Shell, J.W. (1993), World Patent WO 9318755.
- 5 16. Shell, J.W. (1996), U.S. Patent 5,582,837.
17. Fisher, M.C., Morella, A.M. (1994), European Patent 609961.
18. Hansraj, B.R., Bashir, R.H. (1992), European Patent 502642.
- 10 19. Rollet, M. (1987), French Patent 2594693.
20. Howard, S.A., Kotwal, P.M. (1997) U.S. Patent 5,645,858.
21. Macrae, R.J., Smith J.S. (1997), World Patent WO 9718814.
- 15 22. Belenduik, G.W., McCarty, J.A., Rudnic, E.M. (1996), U.S. Patent 5,484,608.
23. Bhatti, G.K., Edgren, D.E., Hatamkhani, Z., Wong, P.S., Wong, P.S.L. (1994), World Patent WO 9427589.
24. Palepu, N.R., Venkatesh, G.M., (1997) European
20 Patent 701436.

What is claimed is:

1. A pharmaceutical formulation comprising (1) an inner solid particulate phase, and (2) an outer solid continuous phase in which particles of the inner solid particulate phase are dispersed and embedded, the particles of the inner solid particulate phase comprising (a) a pharmaceutical having a high water solubility; and (b) an extended release material, and the outer solid continuous phase comprising an extended release material.
2. The pharmaceutical formulation as defined in Claim 1 which is a biphasic heterogeneous controlled release formulation which is designed to release pharmaceutical from the particles forming the inner solid particulate phase through the outer solid continuous phase into the upper gastrointestinal tract.
3. The pharmaceutical formulation as defined in Claim 1 wherein the pharmaceutical is a biguanide antidiabetic agent.
4. The pharmaceutical formulation as defined in Claim 3 wherein the biguanide is metformin or a pharmaceutically acceptable salt thereof.
5. The pharmaceutical formulation as defined in Claim 4 wherein the pharmaceutical is metformin hydrochloride.
6. The pharmaceutical as defined in Claim 1 wherein the inner solid particulate phase is in the form of discrete individual particles or granules and the outer solid continuous phase is a substantially continuous matrix having individual particles forming the inner solid particulate phase embedded therein and dispersed throughout.
7. The pharmaceutical formulation as defined in Claim 1 wherein the pharmaceutical has a solubility in water of at least about 100 mg/ml and a limited window of absorption in the upper gastrointestinal tract.
8. The pharmaceutical formulation as defined in Claim 1 wherein the pharmaceutical present in the inner

solid particulate phase is metformin or a pharmaceutically acceptable salt thereof.

9. The pharmaceutical formulation as defined in Claim 1 wherein the inner solid particulate phase is
5 present in a weight ratio to the outer solid continuous phase within the range from about 0.5:1, to about 4:1.

10. The pharmaceutical formulation as defined in Claim 1 wherein the pharmaceutical is present in the inner solid particulate phase in an amount within the range from
10 about 10 to about 98% by weight of the inner solid particulate phase.

11. The pharmaceutical formulation as defined in Claim 1 wherein the extended release material present in the inner solid particulate phase comprises one or more
15 hydrophilic polymers, one or more hydrophobic polymers and/or one or more other type hydrophobic materials; and the extended release material in the outer solid continuous phase comprises one or more hydrophilic polymers, one or more hydrophobic polymers and/or one or more other type
20 hydrophobic materials.

12. The pharmaceutical formulation as defined in Claim 11 wherein the extended release material present in the inner solid particulate phase comprises one or more ionic polymers and the extended release material present in
25 the outer solid continuous phase comprises one or more non-ionic polymers.

13. The pharmaceutical formulation as defined in Claim 12 wherein the ionic polymer comprises sodium alginate, carbomer, calcium carboxymethylcellulose or
30 sodium carboxymethylcellulose, and the non-ionic polymer comprises hydroxypropylmethylcellulose 2910 USP, viscosity grade ranging from about 4000 to about 100,000 cps and hydroxypropylmethyl cellulose 2208 USP viscosity grade ranging from about 3 to about 150 cps and/or
35 microcrystalline cellulose.

14. The pharmaceutical formulation as defined in Claim 1 wherein the inner solid particulate phase has a

mean particle size within the range from about 30 μm to about 0.8 mm.

15. The pharmaceutical formulation as defined in Claim 1 wherein the inner solid particulate phase comprises metformin, metformin hydrochloride, metformin succinate (2:1) salt or metformin fumarate (2:1) salt, and ethyl cellulose and/or sodium carboxymethyl cellulose and/or glycerylmonostearate and the outer solid continuous phase comprises hydroxypropylmethylcellulose 2208 USP (100,000 cps), and/or hydroxypropylmethylcellulose 2910 USP (5 cps) and/or microcrystalline cellulose.

16. The pharmaceutical formulation as defined in Claim 1 wherein the pharmaceutical is a combination of metformin or a pharmaceutically acceptable salt thereof and another antihyperglycemic agent.

17. A method for preparing a biphasic controlled release delivery system, which comprises forming an inner solid particulate phase comprising individual particles comprising a pharmaceutical having a high water solubility and an extended release material and mixing the individual particles forming the inner solid particulate phase with an outer solid continuous phase comprising an extended release material to thereby disperse and embed the individual particles forming the inner solid particulate phase in the outer solid continuous phase.

18. A biphasic controlled release delivery system formed by the method as defined in Claim 17.

19. A method for treating diabetes which comprises administering to a mammalian patient in need of treatment a therapeutically effective amount of the formulation as defined in Claim 4.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US99/05233

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) :A61K 9/24

US CL :424/457

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 424/472

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 5,645,858 A (KOTWAL ET AL.) 08 July 1997, see cols. 5-8.	1-19
Y	WO 96/08243 A1 (MOECKEL ET AL.) 21 March 1996, see entire disclosure.	1-19
Y	EP 0 609 961 A1 (MORELLA ET AL.) 10 August 1994, see examples and tables.	1-19

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
B earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*A* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

03 JUNE 1999

Date of mailing of the international search report

07 JUL 1999

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

PETER KULKOSKY *For*

Telephone No. (703) 308-0196